## Biology 1200

Organic molecules – nucleotides Nucleic Acids

Organic Molecules: Nucleotides

- 3 principle components
  - nitrogenous base
    - single or double carbon-nitrogen ring
  - sugar (monosaccharide)
  - · one or more phosphate groups
- ATP contains • adenine


2

- 3 phosphate groups
- ATP is the universal energy carrying molecule

1

ATP • High energy bonds

- second and third phosphate groups are attached by high energy covalent bonds
- phosphate groups are negatively charged and naturally repel each other
- ATPases hydrolyze the  $3^{\rm rd}$  high energy phosphate bond of ATP producing ADP +  $P_i$  + energy
- Kinases (phosphokinases)
  - enzymes that phosphorylate (add the Pireleased from ATP to) other enzymes or molecules to activate them



3

Production and Uses of ATP Glucose + 6 O2 are converted to + 6 CO2 + 6 H2O ADP + Pi

ATP Production - Glycolysis • Glycolysis splits one 6 carbon glucose into two 3 carbon pyruvic acid molecules • yield 2 net ATP's

6 5


1

## ATP Production - Anaerobic Fermentation

- If no oxygen is available pyruvic acid is converted to lactic acid (build up causes muscle soreness)
- Some ATP produced


7

• Allows glycolysis to start over (regenerates NAD+)



## ATP Production - Aerobic Respiration If oxygen is available pyruvic acid is efficiently consumed yielding 36 more ATP molecules (from the original glucose) Oxygen • Aerobic respiration occurs in mitochondrion CO2 + H2O 36 ADP + 36 Pi

8



Other Nucleotides

- Guanosine triphosphate (GTP)
  - may donate a phosphate group (Pi) to other molecules or to ADP
- Cyclic adenosine monophosphate (cAMP) formed after removal of both high energy Pi's

  - after chemical signal (first messenger) binds to cell surface, it triggers the conversion of ATP to cAMP (second messenger) to activate effects inside cell
- Nucleic acids are polymers of nucleotides

9

10

**Nucleic Acids** 

- DNA (deoxyribonucleic acid)
  - 100 million to 1 billion nucleotides long
  - contains the genetic code for
    - cell division, sexual reproduction, the instructions for protein synthesis
- RNA (ribonucleic acid)
  - 3 forms of RNA range from 70 to 10,000 nucleotides long

  - carries out instructions given by DNA
     synthesizes the proteins coded for by DNA

12 11

## The Instructions for Life Nucleic Acids

- Proteins which control or make a vast majority of our body are made from codes written on genes. Genes are parts of DNA that contain nucleotides in a particular order.
- There are two types of nucleic acids: ribonucleic acid (RNA) and deoxyribonucleic acid (DNA).
- DNA gives direction for its own replication.
- DNA also directs RNA synthesis and, through RNA, controls protein synthesis.
- Organisms inherit DNA from their parents.
  - Each DNA molecule is extremely long and typically consists of hundreds to thousands of genes.
  - When a cell reproduces itself by mitosis or division, its DNA is copied and passed to the next generation of cells.

**Nucleic Acids** 

- · Nucleic acids are many nucleotides joined together.
- Each nucleotide consists of three parts: a nitrogen base, a pentose, 5 carbon, sugar, and a phosphate group

- and a phosphate group

  Purines and pyrimidines are the 2 types of nucleotides.


  Pyrimidines have a single six-membered ring.


  The three different pyrimidines, cytosine (C), thymine (T), and uracil (U)—part of RNA.

  Purine have a six-membered ring joined to a five-membered ring. So have 2 rings not 1

  The two purines are adenine (A) and guanine (G).

13 14





15 16